

# A Note on the Erdős–Szemerédi LCM-Window Problem for Exponents Greater Than $1/4$

Repaired version, 30 April 2026

## Abstract

Let  $A = \{a_1 < a_2 < \dots\} \subseteq \mathbb{N}$ , and let  $F(A, X, k)$  denote the number of indices  $i$  for which

$$\text{lcm}(a_i, a_{i+1}, \dots, a_{i+k-1}) < X.$$

Using a recent divisor-in-short-interval estimate of Letendre, we prove that for every  $\varepsilon > 1/4$  there is a fixed  $k = k(\varepsilon)$  such that

$$F(A, X, k) \ll_{\varepsilon, k} X^\varepsilon$$

uniformly in  $A$ .

## 1 Input from divisors in short intervals

For an integer  $n \geq 1$ , write

$$D_n(U, H) = \#\{d \mid n : U \leq d \leq U + H\}.$$

We use the following proposition of Letendre [1, Proposition 1].

**Proposition 1** (Letendre). *Let  $n \geq 1$  be fixed, and let  $0 < \theta < 1$  and  $0 < \eta < \theta^2$ . Then*

$$D_n(n^\theta, n^{\theta^2-\eta}) \ll \frac{\theta(1-\theta)}{\eta} + \frac{1}{\theta(1-\theta)}.$$

*In particular, for every  $0 < \lambda < 1/2$  and every  $0 < \eta < \lambda^2$ , there is a constant  $C(\lambda, \eta)$  such that*

$$D_n(n^\theta, n^{\theta^2-\eta}) \leq C(\lambda, \eta)$$

*for all  $n \geq 1$  and all  $\lambda \leq \theta \leq 1 - \lambda$ .*

## 2 Two elementary packing lemmas

For a good index  $i$ , put

$$L_i = \text{lcm}(a_i, a_{i+1}, \dots, a_{i+k-1}), \quad u_i = a_i, \quad H_i = a_{i+k-1} - a_i.$$

Then  $L_i < X$ , and the interval  $[u_i, u_i + H_i]$  contains the  $k$  divisors

$$a_i, a_{i+1}, \dots, a_{i+k-1}$$

of  $L_i$ .

**Lemma 1** (Disjointness after passing to one residue class). *Fix  $k \geq 1$  and one residue class modulo  $k$ . The intervals*

$$[a_i, a_{i+k-1}]$$

*attached to good indices  $i$  in this residue class are pairwise disjoint.*

*Proof.* If  $i < j$  and  $i \equiv j \pmod{k}$ , then  $j \geq i + k$ . Since the sequence  $A$  is strictly increasing,

$$a_{i+k-1} < a_j.$$

Thus the interval attached to  $i$  lies strictly to the left of the interval attached to  $j$ .  $\square$

**Lemma 2** (Packing by separated left endpoints). *Let  $\mathcal{I}$  be a finite family of pairwise disjoint intervals  $[u, u + H]$  with integer left endpoints  $u \in [U, 2U)$ . If every interval in  $\mathcal{I}$  has length at least  $h > 0$ , then*

$$\#\mathcal{I} \ll \frac{U}{h} + 1,$$

*with an absolute implied constant.*

*Proof.* Order the intervals by increasing left endpoint. If two consecutive intervals have left endpoints  $u < v$ , then disjointness and the lower bound on the first length give

$$v \geq u + H \geq u + h,$$

up to an immaterial endpoint convention. Hence the left endpoints are  $h$ -separated inside an interval of length  $U$ , and there are  $O(U/h + 1)$  of them.  $\square$

### 3 Proof of the $\varepsilon > 1/4$ bound

**Theorem 1.** *For every  $\varepsilon > 1/4$ , there exists an integer  $k = k(\varepsilon)$  such that, uniformly for every increasing sequence  $A \subseteq \mathbb{N}$ ,*

$$F(A, X, k) \ll_{\varepsilon, k} X^\varepsilon.$$

*Consequently, for every  $\varepsilon > 1/4$ , the Erdős–Szemerédi inequality*

$$F(A, X, k) < X^\varepsilon$$

*holds for all sufficiently large  $X$ , with  $k$  depending only on  $\varepsilon$ .*

*Proof.* Fix  $\varepsilon > 1/4$ . Choose  $\lambda$  and  $\eta$  such that

$$0 < \lambda < \frac{1}{2}, \quad \frac{\lambda}{1-\lambda} < \varepsilon, \quad 0 < \eta < \min\{\lambda^2, \varepsilon - 1/4\}.$$

Let  $C(\lambda, \eta)$  be the uniform constant from Proposition 1, and choose an integer

$$k > \max\{2, C(\lambda, \eta) + 1\}.$$

We shall prove that this  $k$  works.

By Lemma 1, it is enough to bound the number of good indices in a fixed residue class modulo  $k$  and then multiply by  $k$ . Hence, throughout the proof, restrict to one residue class modulo  $k$ ; all associated intervals  $[u_i, u_i + H_i]$  are pairwise disjoint.

For a good index  $i$ , define

$$L_i = \text{lcm}(a_i, a_{i+1}, \dots, a_{i+k-1}) < X.$$

Since  $u_i = a_i$  divides  $L_i$ , and since  $u_i \leq L_i$ , we may write

$$u_i = L_i^{\theta_i}$$

for some  $0 \leq \theta_i \leq 1$ , except for harmless cases with  $u_i = 1$  or  $L_i = 1$ . These exceptional cases contribute only  $O_k(1)$  indices:  $u_i = 1$  can occur for at most one index, and  $L_i = 1$  is impossible for  $k \geq 2$  in a strictly increasing sequence of positive integers.

We split the remaining good indices into three classes.

### 1. Small $\theta_i$

Suppose  $\theta_i < \lambda$ . Then

$$u_i = L_i^{\theta_i} < X^\lambda.$$

There are at most  $O(X^\lambda)$  possible integer values of  $u_i$ , so this contribution is  $O(X^\lambda) = O(X^\varepsilon)$ .

### 2. Large $\theta_i$

Suppose  $\theta_i > 1 - \lambda$ . We use a simple spacing estimate for divisors near the top of  $L_i$ . If  $d < e$  are two distinct divisors of  $L_i$ , write  $d = L_i/r$  and  $e = L_i/s$  with integers  $r > s \geq 1$ . Then

$$e - d = L_i \left( \frac{1}{s} - \frac{1}{r} \right) = \frac{L_i(r - s)}{rs} \geq \frac{L_i}{r^2} = \frac{d^2}{L_i}.$$

In our interval all listed divisors are at least  $u_i = L_i^{\theta_i}$ , so consecutive listed divisors in the interval are separated by at least

$$\frac{u_i^2}{L_i} = u_i^{2-1/\theta_i} \geq u_i^{(1-2\lambda)/(1-\lambda)},$$

because  $\theta_i > 1 - \lambda$  and the function  $2 - 1/t$  is increasing for  $t > 0$ .

Group intervals by dyadic ranges  $U \leq u_i < 2U$ . In such a range, every interval of the large- $\theta_i$  type has length

$$H_i \gg_k U^{(1-2\lambda)/(1-\lambda)},$$

because it contains  $k$  listed divisors and hence  $k - 1$  gaps of the above size. By Lemma 2, the number of such intervals in this dyadic range is

$$O_k \left( \frac{U}{U^{(1-2\lambda)/(1-\lambda)}} + 1 \right) = O_k \left( U^{\lambda/(1-\lambda)} + 1 \right).$$

Summing over dyadic  $1 \leq U < X$  gives

$$O_k \left( X^{\lambda/(1-\lambda)} \log X \right) = O_{\varepsilon,k}(X^\varepsilon),$$

because  $\lambda/(1 - \lambda) < \varepsilon$ .

### 3. Middle range $\lambda \leq \theta_i \leq 1 - \lambda$

It remains to treat good indices with

$$\lambda \leq \theta_i \leq 1 - \lambda.$$

We use the uniform compact-range consequence of Proposition 1. Since  $k > C(\lambda, \eta)$ , the interval  $[u_i, u_i + H_i]$  cannot have length less than  $L_i^{\theta_i^2 - \eta}$ . Indeed, if it had smaller length, then the larger divisor-counting interval

$$[L_i^{\theta_i}, L_i^{\theta_i} + L_i^{\theta_i^2 - \eta}]$$

would contain the same  $k$  listed divisors of  $L_i$ , contradicting

$$D_{L_i}(L_i^{\theta_i}, L_i^{\theta_i^2 - \eta}) \leq C(\lambda, \eta) < k.$$

Thus

$$H_i \geq L_i^{\theta_i^2 - \eta}.$$

Since  $u_i = L_i^{\theta_i}$ , this gives

$$H_i \geq u_i^{\theta_i} L_i^{-\eta} > u_i^{\theta_i} X^{-\eta}.$$

Again group indices dyadically, say

$$U \leq u_i < 2U, \quad \alpha = \frac{\log U}{\log X}, \quad 0 \leq \alpha \leq 1.$$

Because  $u_i \geq U$  and  $L_i < X$ , we have

$$\theta_i = \frac{\log u_i}{\log L_i} \geq \frac{\log U}{\log X} = \alpha.$$

Therefore

$$u_i^{\theta_i} \geq U^{\theta_i} \geq U^\alpha = X^{\alpha^2},$$

and hence

$$H_i \geq X^{\alpha^2 - \eta}.$$

By Lemma 2, the number of middle-range intervals with  $U \leq u_i < 2U$  is

$$O\left(\frac{U}{X^{\alpha^2 - \eta}} + 1\right) = O\left(X^{\alpha - \alpha^2 + \eta} + 1\right).$$

Since

$$\alpha - \alpha^2 \leq \frac{1}{4}$$

for all real  $\alpha$ , this is

$$O\left(X^{1/4 + \eta} + 1\right).$$

Summing over the  $O(\log X)$  dyadic choices of  $U$  yields

$$O\left(X^{1/4 + \eta} \log X\right) = O_\varepsilon(X^\varepsilon),$$

because  $1/4 + \eta < \varepsilon$ .

Combining the three cases, the number of good indices in the chosen residue class is  $O_{\varepsilon,k}(X^\varepsilon)$ . Multiplying by the  $k$  residue classes modulo  $k$  gives

$$F(A, X, k) \ll_{\varepsilon,k} X^\varepsilon,$$

as claimed.

For the final displayed inequality with the strict exponent  $\varepsilon$ , apply the just-proved big- $O$  statement with any exponent  $\varepsilon'$  satisfying  $1/4 < \varepsilon' < \varepsilon$ . Since the implied constant is uniform in  $A$ , we have  $C_{\varepsilon',k} X^{\varepsilon'} < X^\varepsilon$  for all sufficiently large  $X$ .  $\square$

**Remark 1.** *The endpoint  $\varepsilon = 1/4$  is not obtained by this argument. The exponent  $1/4$  appears from the elementary maximum*

$$\max_{0 \leq \alpha \leq 1} (\alpha - \alpha^2) = \frac{1}{4}.$$

*The example  $A = \mathbb{N}$  also gives the lower obstruction  $F(\mathbb{N}, X, k) \gg_k X^{1/k}$ . Thus the proof above reaches every exponent strictly larger than  $1/4$ , but not the endpoint.*

## References

- [1] Patrick Letendre, *Divisors of an Integer in a Short Interval*, arXiv:2503.12146, 2025.